arXiv:1608.03915v2 [math.NT] 4 Sep 2016

The action of $\mathrm{GL}_2(\mathbb{F}_q)$ on irreducible polynomials over $\mathbb{F}_q$

Lucas Reis

Departamento de Matemática, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte, MG, 30123-970, Brazil

Abstract

Let $\mathbb{F}_q$ be the finite field with q elements, $p = \mathrm{char}\mathbb{F}_q$. The group $\mathrm{GL}_2(\mathbb{F}_q)$ acts naturally in the set of irreducible polynomials over $\mathbb{F}_q$ of degree at least 2. In this paper we are interested in the characterization and number of the irreducible polynomials that are fixed by the elements of a subgroup H of $\mathrm{GL}_2(\mathbb{F}_q)$. We make a complete characterization of the fixed polynomials in the case when H has only elements of the form $\begin{pmatrix}1 & b\\ 0 & 1\end{pmatrix}$, corresponding to translations $x \mapsto x + b$ and, as a consequence, the case when H is a $p-$subgroup of $\mathrm{GL}_2(\mathbb{F}_q)$. This paper also contains alternative solutions for the cases when H is generated by an element of the form $\begin{pmatrix}a & 0\\ 0 & 1\end{pmatrix}$, obtained by Garefalakis (2010) [3] and $H = \mathrm{PGL}_2(\mathbb{F}_q)$, obtained by Stichtenoth and Topuzoglu (2011) [6].

Keywords: finite fields, irreducible polynomials, group action
2010 MSC: 12E20, 11T55

1. Introduction

Let $\mathbb{F}_q$ be the finite field with q elements, $p = \mathrm{char}\mathbb{F}_q$. As it was noticed in [3] and [6], there is a natural action of the group $\mathrm{GL}_2(\mathbb{F}_q)$ on the set I of irreducible polynomials of degree at leat 2 in $\mathbb{F}_q[x]$. Namely, given $A = \begin{pmatrix}a & b\\ c & d\end{pmatrix}$ in $\mathrm{GL}_2(\mathbb{F}_q)$ and $f(x) \in I$ of degree n we can define

$$A \circ f = (cx + d)^n f\left(\frac{ax + b}{cx + d}\right).$$

It can be verified that f and $A \circ f$ have the same degree and $A \circ (B \circ f) = (AB) \circ f$ for all $A, B \in \mathrm{GL}_2(\mathbb{F}_q)$. From this definition, two interesting theoretical questions arise:

a) Given a subgroup H of $\mathrm{GL}_2(\mathbb{F}_q)$, which elements $f \in I$ are fixed by H, i.e., $A \circ f = f$ for all $A \in H$?

b) How many fixed elements exist?

In [6], the authors gave a complete characterization of the elements $f \in I$ that are fixed by $H = \langle A \rangle$, where A is any element of $\mathrm{GL}_2(\mathbb{F}_q)$. Earlier, the same characterization was given in [3] for the special cases $A = \begin{pmatrix}a & 0\\ 0 & 1\end{pmatrix}$ or $\begin{pmatrix}1 & b\\ 0 & 1\end{pmatrix}$, corresponding to the homothety $x \mapsto ax$ and the translation $x \mapsto x + b$,

Email address: lucasreismat@gmail.com (Lucas Reis)

respectively, including enumeration formulas. Combining those results, in [3] the author also obtained enumeration formulas for the case when A is an upper triangular matrix in $\mathrm{GL}_2(\mathbb{F}_q)$.

In this paper we are also going to consider only translations and homotheties but with an alternative characterization of the invariant polynomials. Section 2 is devoted to extend the results of Garefalakis [3] for translations, considering now any family of translations $\{x \mapsto x + s; s \in S\}$ where $S \subset \mathbb{F}_q$. Section 3 includes alternative proofs of the enumeration formula in [3] for homotheties and the characterization of all irreducible polynomials that are invariant under the action of the whole group $\mathrm{PGL}_2(\mathbb{F}_q)$, [[6], Proposition 4.8]. In Section 4, using the canonical conjugation in $\mathrm{GL}_2(\mathbb{F}_q)$ and the results of Section 2, we count the irreducible polynomials that are fixed by any given $p-$subgroup H of $\mathrm{GL}_2(\mathbb{F}_q)$.

2. S-Translation Invariant Polynomials

Throughout this section $\mathbb{F}_q$ denotes the finite field with q elements, where $q = p^k$ is a prime power. If $S \subset \mathbb{F}_q$, we say that $f(x) \in \mathbb{F}_q[x]$ is an S-translation invariant polynomial if $f(x + s) = f(x)$ for all $s \in S$.

Example 1. *$f(x) = x^q - x$ is an $\mathbb{F}_q$-translation invariant polynomial.*

Let $C_S(n)$ denote the set of all S-translation invariant monic irreducible polynomials of degree n over $\mathbb{F}_q$. Suppose that $a, b \in \mathbb{F}_q^*$ and $f(x) \in \mathbb{F}_q[x]$ is a polynomial such that $f(x + a) = f(x)$ and $f(x + b) = f(x)$. Clearly $f(x + 0) = f(x)$ and then, by induction, we have that

$$f(x + ia + jb) = f(x)$$

for all $i, j \in \mathbb{F}_p$. In particular, if $S' \subset \mathbb{F}_q$ is the $\mathbb{F}_p-$vector space generated by S, then $C_{S'}(n) \supset C_S(n)$. The converse inclusion is obviously true and so $C_{S'}(n) = C_S(n)$. From now, $S \subset \mathbb{F}_q$ will denote an $\mathbb{F}_p-$vector space of dimension $r > 0$.

Remark 1. *If $f(x) \in \mathbb{F}_q[x]$ is an S-translation invariant polynomial and $a \in \mathbb{F}_q^*$, then $g(x) = a^n f(a^{-1}x)$ is an S'-translation invariant polynomial, where $S' = \{as | s \in S\}$. Moreover, if S is an $\mathbb{F}_p-$vector space then so is S'.*

The observations above lead us to the following definition:

Definition 1. *Let $S, S' \subset \mathbb{F}_q$ be $\mathbb{F}_p$-vector spaces. We say that S and S' are $\mathbb{F}_q-$linearly equivalent and write $S \sim_{\mathbb{F}_q} S'$ if there exists $a \in \mathbb{F}_q^*$ such that $S' = \{as | s \in S\}$.*

It is immediate from definition that the relation $\sim_{\mathbb{F}_q}$ is an equivalence relation. Moreover, this relation gives an interesting invariant:

Lemma 1. *If $S \sim_{\mathbb{F}_q} S'$, then $|C_S(n)| = |C_{S'}(n)|$ for all $n \in \mathbb{N}$.*

Proof. If $S \sim_{\mathbb{F}_q} S'$, then $S' = \{as | s \in S\}$ for some $a \in \mathbb{F}_q^*$. Let $I_n(q)$ be the set of all monic irreducible polynomials of degree n over $\mathbb{F}_q$ and

$$\begin{array}{rccc} \tau_a : & I_n(q) & \to & I_n(q) \\ & f(x) & \mapsto & a^n f(a^{-1}x). \end{array}$$

Clearly τ_a is well defined and is one to one. From Remark 1 we know that if $f(x) \in C_S(n)$ then $\tau_a(f(x)) \in C_{S'}(n)$. Therefore we have that $|C_S(n)| \leq |C_{S'}(n)|$. Since the relation $\sim_{\mathbb{F}_q}$ is symmetric, it follows that $|C_{S'}(n)| \leq |C_S(n)|$ and this completes the proof. □

The following theorem gives a characterization of the S-translation invariant polynomials over the finite field $\mathbb{F}_q$.

Theorem 1. *Let $S \subset \mathbb{F}_q$ be an $\mathbb{F}_p-$vector space of dimension $r > 0$ and*

$$P_S(x) := \prod_{s \in S} (x - s).$$

Then $g(x) \in \mathbb{F}_q[x]$ is an S-translation invariant polynomial if, and only if, there exists some polynomial $f(x) \in \mathbb{F}_q[x]$ such that $g(x) = f(P_S(x))$. In particular, any S-translation invariant polynomial has degree divisible by $\deg P_S(x) = |S| = p^r$.

Proof. Assume that $g(x)$ is an S-translation invariant polynomial over $\mathbb{F}_q$. We proceed by induction on $n = \deg g(x)$. If $g(x)$ is constant then there is nothing to prove. Suppose that the statement is true for all polynomials of degree at most k and let $g(x)$ be an S-translation invariant polynomial of degree $k + 1$. We have $g(0) = g(s)$ for all $s \in S$ and so the polynomial $g(x) - g(0)$ has degree $k + 1 > 0$ and vanishes at s for all $s \in S$. In particular we have that

$$g(x) - g(0) = P_S(x)G(x) \tag{1}$$

for some non-zero polynomial $G(x) \in \mathbb{F}_q[x]$. Since $g(x+s) - g(0) = g(x) - g(0)$ and $P_S(x+s) = P_S(x)$ for all $s \in S$, by equation (1) it follows that $G(x)$ is an S-translation invariant polynomial over $\mathbb{F}_q$ and $\deg G(x) < \deg g(x)$. By the induction hypothesis we have that $G(x) = F(P_S(x))$ for some $F(x) \in \mathbb{F}_q[x]$. Therefore $g(x) = P_S(x)F(P_S(x)) + g(0)$ and so $g(x) = f(P_S(x))$ where $f(x) = xF(x) + g(0)$. The converse is obviously true. □

If $1, a \in S$ where $a \notin \mathbb{F}_p$, we have the following:

Corollary 1. *Let $S \subset \mathbb{F}_q$ be an $\mathbb{F}_p-$vector space such that $1, a \in S$ where $a \notin \mathbb{F}_p$. Then any S-translation invariant polynomial in $\mathbb{F}_q[x]$ is of the form*

$$f(x^{p^2} - x^p(1 + (a - a^p)^{p-1}) + x(a - a^p)^{p-1})$$

for some polynomial $f(x) \in \mathbb{F}_q[x]$.

Proof. Since $1, a \in S$ and $a \notin \mathbb{F}_p$, we have that any S-translation invariant polynomial is also a S'-translation invariant where $S' = \langle 1, a \rangle_{\mathbb{F}_p} = \{j + ai | i, j \in \mathbb{F}_p\}$. In addition, notice that

$$P_{S'}(x) = \prod_{0 \le i,j \le p-1} (x - j - ai) = x^{p^2} - x^p(1 + (a - a^p)^{p-1}) + x(a - a^p)^{p-1}.$$

□

The main result of this paper is the following:

Theorem 2. *Let $\mathbb{F}_q$ be the finite field with $q = p^k$ elements and $S \in \mathbb{F}_q$ be an $\mathbb{F}_p-$vector space of dimension $r > 0$.*

a) If $r > 1$ then $|C_S(n)| = 0$ for all n.

b) If $r = 1$ and n is not divisible by p then $|C_S(n)| = 0$.

c) If $r = 1$ and $n = pm$, then

$$|C_S(n)| = \frac{p-1}{pm} \sum_{\substack{d|m \\ \gcd(d,p)=1}} q^{m/d} \mu(d).$$

2.1. Lemmata

In order to prove Theorem 2 we have to discuss the irreducibility of the polynomials $f(P_S(x)) \in \mathbb{F}_q[x]$. In this direction, Lemmas 2 and 3 give some criteria to ensure the irreducibility of special polynomial compositions and Lemma 4 will be useful in the proof of the enumeration formula in Theorem 2.

Lemma 2 ([1])**.** *Let $f(x) = x^n + Bx^{n-1} + \cdots + c \in \mathbb{F}_q[x]$ be an irreducible polynomial, where $q = p^k$ is a prime power. The polynomial $f(x^{p^{2t}} - ax^{p^t} - bx)$ is also irreducible over $\mathbb{F}_q[x]$ if and only if the following conditions are satisfied:*

a) $p = 2, t = 1, n$ *is odd and* $B \neq 0$,

b) $\gcd(x^3 - ax - b, x^{2^k} - x) \neq 1$,

c) $\mathrm{Tr}_{L/K}(\beta^{-2}B) = \mathrm{Tr}_{L/K}(\alpha^{-2}\beta) = 1$, *where* $L = \mathbb{F}_{2^k}, K = \mathbb{F}_2$ *and* α, β *are two elements in* L *such that* $\alpha^2 + \beta = a$ *and* $\alpha\beta = b$.

Lemma 3 ([4], Theorem 3.82)**.** *Let $f(x) = x^n + a_{n-1}x^{n-1} + \cdots + c \in \mathbb{F}_q[x]$ be an irreducible polynomial, $p = \mathrm{char}\mathbb{F}_q$ and $b \in \mathbb{F}_q$. The polynomial $f(x^p - x - b)$ is also irreducible over $\mathbb{F}_q[x]$ if and only if $\mathrm{Tr}_{L/K}(nb - a_{n-1}) \neq 0$ where $L = \mathbb{F}_q$ and $K = \mathbb{F}_p$.*

Lemma 4 ([5])**.** *The number of monic irreducible polynomials over $\mathbb{F}_q$ with degree n and a given trace $a \neq 0$ is*

$$\frac{1}{qn} \sum_{\substack{d|n \\ \gcd(d,p)=1}} q^{n/d}\mu(d).$$

2.2. Proof of Theorem 2

If $S \subset \mathbb{F}_q$ is any $\mathbb{F}_p-$vector space of dimension $r > 0$ and $a \in S \setminus \{0\}$, then $a^{-1}S \sim_{\mathbb{F}_q} S$ and $1 \in a^{-1}S$. Thus, using Lemma 1, we can suppose that $1 \in S$.

a) Let n be a positive integer and $S \subset \mathbb{F}_q$ be an $\mathbb{F}_p-$vector space of dimension $r > 1$ such that $1 \in S$. Since $r > 1$, there is some element $\gamma \in S \setminus \mathbb{F}_p$. It follows from Corollary 1 that any $g(x) \in C_S(n)$ is of the form

$$f(x^{p^2} - x^p(1 + (\gamma - \gamma^p)^{p-1}) + x(\gamma - \gamma^p)^{p-1}).$$

Therefore, by Lemma 2, $g(x)$ is reducible whenever $p > 2$.

If $p = 2$, we have that $g(x) = f(x^4 - ax^2 - bx)$ where $a = \gamma^2 + \gamma + 1$ and $b = \gamma^2 + \gamma$. Now consider the following system of equations:

$$\begin{cases} \alpha^2 + \beta &= a \\ \alpha\beta &= b. \end{cases} \tag{2}$$

Notice that, for any solution (α, β) of system (2), α is a root of the equation $y^3 - ay + b = 0$ and $\beta = a - \alpha^2$. In particular the system (2) has at most three solutions and since $(\alpha, \beta) = (1, \gamma^2 + \gamma), (\gamma, \gamma + 1), (\gamma + 1, \gamma)$ are solutions these are all of them.

Condition c) of Lemma 2 says that, for some solution (α,β) of (2), we have $\mathrm{Tr}_{K/L}(\alpha^{-2}\beta) = 1$ where $L = \mathbb{F}_{2^k}$ and $K = \mathbb{F}_2$. In particular, if $(\alpha,\beta) = (1, \gamma^2+\gamma)$ then

$$\mathrm{Tr}_{K/L}(\alpha^{-2}\beta) = \mathrm{Tr}_{K/L}(\gamma^2) + \mathrm{Tr}_{K/L}(\gamma) = \mathrm{Tr}_{K/L}(\gamma) + \mathrm{Tr}_{K/L}(\gamma) = 0 \neq 1.$$

The other two solutions of (2) satisfy $\beta = \alpha + 1$ and in these cases we have

$$\mathrm{Tr}_{K/L}(\alpha^{-2}\beta) = \mathrm{Tr}_{K/L}(\alpha^{-1}) + \mathrm{Tr}_{K/L}((\alpha^{-1})^2) = \mathrm{Tr}_{K/L}(\alpha^{-1}) + \mathrm{Tr}_{K/L}(\alpha^{-1}) = 0 \neq 1.$$

Thus $g(x)$ is never irreducible if $r > 1$ and this completes the proof of a).

b) This follows directly from Theorem 1 since any polynomial of the form $f(P_S(x))$ has degree divisible by $|S| = p$.

c) Since $r = 1$ and $1 \in S$, we have $S = \mathbb{F}_p$. Let m be any positive integer and $g(x)$ be an irreducible polynomial of degree pm. From Lemma 1 we have that $g(x) \in C_S(pm)$ if, and only if, $g(x)$ is of the form $f(P_S(x)) = f(x^p - x)$ where $f(x) = x^m + a_{m-1}x^{m-1} + \cdots + a_0$ is an irreducible polynomial over $\mathbb{F}_q$.

From Lemma 3, the polynomial $f(x^p - x) \in \mathbb{F}_q[x]$ is irreducible if, and only if, $\mathrm{Tr}_{L/K}(a_{m-1}) \neq 0$ where $L = \mathbb{F}_q$ and $K = \mathbb{F}_p$. It is well known that for each $a \in K$, the equation $\mathrm{Tr}_{L/K}(x) = a$ has $\frac{q}{p}$ distinct solutions. Thus for exactly $(p-1)\frac{q}{p}$ values of $a_{m-1} \in \mathbb{F}_q$ we have $\mathrm{Tr}_{L/K}(-a_{m-1}) \neq 0$ and any of these elements is nonzero. For a fixed $\beta \in \mathbb{F}_q^*$, by Lemma 4, there exist exactly

$$\frac{1}{qm} \sum_{\substack{d|m \\ \gcd(d,p)=1}} q^{m/d}\mu(d)$$

monic irreducible polynomials over $\mathbb{F}_q$ with degree m and trace β. Thus the number of monic irreducible polynomials $f(x) \in \mathbb{F}_q[x]$ of degree m such that $f(x^p - x) \in \mathbb{F}_q[x]$ is also irreducible is equal to

$$(p-1)\frac{q}{p} \cdot \frac{1}{qm} \sum_{\substack{d|m \\ \gcd(d,p)=1}} q^{m/d}\mu(d) = \frac{p-1}{pm} \sum_{\substack{d|m \\ \gcd(d,p)=1}} q^{m/d}\mu(d).$$

Since the polynomials $f(x^p - x)$ are all distinct when $f(x)$ runs through $I_q(m)$, we are done.

3. Miscellanea

Using the results obtained in Section 2, we give alternative proofs of two interesting results concerning the action of $\mathrm{GL}_2(\mathbb{F}_q)$ on irreducible polynomials.

3.1. Polynomials invariant under homotheties

Given $a \in \mathbb{F}_q \setminus \{0,1\}$, we are interested in counting the number of monic irreducible polynomials $g(x)$ that are invariant under the homothety $x \mapsto ax$, i.e., $g(ax) = g(x)$. We have the following characterization:

Theorem 3. *Let $f(x)$ be a polynomial over $\mathbb{F}_q$. Then $f(ax) = f(x)$ if and only if there exists $g(x) \in \mathbb{F}_q[x]$ such that*

$$f(x) = g(P_a(x)),$$

where $P_a(x) = \prod_{i=0}^{k-1}(x - a^i) = x^k - 1$. In particular any polynomial invariant under the homothety $x \mapsto ax$ has degree divisible by $k = \mathrm{ord}(a)$.

Proof. Notice that if $f(ax) = f(x)$ and $k = \mathrm{ord}(a)$ is the order of $a \in \mathbb{F}_q$, then $f(1) = f(a^i)$ for all $0 \leq i \leq k-1$; from now, the proof is quite similar to the one of Theorem 1. □

Let $N_a(nk)$ be the number of monic irreducible polynomials $f(x) \in \mathbb{F}_q[x]$ of degree nk such that $f(ax) = f(x)$, $k = \mathrm{ord}(a)$. Also, let $L(n,k)$ be the number of monic irreducible polynomials of the form $F(x^k)$, where F has degree n. Notice that, from Theorem 3, $N_a(nk)$ is exactly the number of monic irreducible polynomials of the form $f(x^k - 1)$ where f has degree n. Since $f(x^k - 1) = F(x^k)$ for $F(x) = f(x-1)$ and $\{f(x-1) | f \in I_q(n)\} = I_q(n)$, the number of monic irreducible polynomials F of degree n for which $F(x^k)$ is also irreducible is the same for the composition $F(x^k - 1)$. In particular we have proved that $N_a(nk) = L(n,k)$. Combining the previous equality with the enumeration formula for $L(n,k)$ presented in [[2], Theorem 3] we directly deduce the enumeration formula of Garefalakis for homotheties:

Theorem 4 ([3], Theorem 4)**.** *If n is not divisible by $k = \mathrm{ord}(a)$, then $N_a(n) = 0$ and, if $n = mk$, we have*

$$N_a(mk) = \frac{\varphi(k)}{mk} \sum_{\substack{d|m \\ \gcd(d,k)=1}} \mu(d)(q^{m/d} - 1).$$

3.2. The action of $\mathrm{PGL}_2(\mathbb{F}_q)$ *on irreducible polynomials*

Consider the projective linear group $G = \mathrm{PGL}_2(\mathbb{F}_q) \approx \mathrm{GL}_2(\mathbb{F}_q)/\sim$, where $A \sim B$ if $A = \lambda B$ for some $\lambda \in \mathbb{F}_q^*$. We are interested to find the monic irreducible polynomials $f(x) \in \mathbb{F}_q[x]$ such that $A \circ f = f$ for any $A \in G$. As it was shown in [[6], Proposition 4.8] in general there are no such polynomials:

Theorem 5. *Let $f \in \mathbb{F}_q[x]$ be a monic irreducible polynomial of degree $n \geq 2$. Suppose that $A \circ f = f$ for all $A \in \mathrm{PGL}_2(\mathbb{F}_q)$. Then $n = q = 2$ and $f(x) = x^2 + x + 1 \in \mathbb{F}_2[x]$.*

Here we give an alternative proof of this fact: let $f \in \mathbb{F}_q[x]$ be as above. Since all 2×2 matrices corresponding to translations belong to $\mathrm{PGL}_2(\mathbb{F}_q)$, f must be an $\mathbb{F}_q-$translation invariant and, from Theorem 2 part a), we know that this is only possible when $\mathbb{F}_q$ is a prime field. Thus $q = p$ prime. Also, from Theorem 1, we know that there exists some polynomial $g(x)$ with nonzero trace such that $f(x) = g(x^p - x)$.

Now, if $p > 2$, then there is some element $a \in \mathbb{F}_p \setminus \{0,1\}$ such that $k = \mathrm{ord}(a) > 1$. Since all homotheties also belong to $\mathrm{PGL}_2(\mathbb{F}_q)$, it follows that $f(x) = f(ax)$ or, equivalently, $g(x^p - x) = g_2(x^p - x)$, where $g_2(x) = g(ax)$. Therefore $g(x) = g_2(x)$, i.e., $g(x)$ is also invariant under the homothety $x \mapsto ax$. From Theorem 3 we have that $g(x) = h(x^k - 1)$ for some $h(x) \in \mathbb{F}_p[x]$. Since $k > 1$, a direct calculation shows that $g(x)$ has trace equal to zero, a contradiction.

Thus $p = 2$, $f(x) = g(x^2 + x)$ and $\deg f(x) = 2N = 2 \deg g(x)$. Let

$$\gamma, \gamma^2, \cdots, \gamma^{2^{2N-1}}$$

be the roots of $f(x)$. Notice that $f(\gamma+1) = g(\gamma^2+\gamma) = 0$ and then $\gamma + 1 = \gamma^{2^j}$ for some $0 < j < 2N$. In other words, $F(x) = x^{2^j} + x + 1 \in \mathbb{F}_2[x]$ has γ as a root. By hypothesis, $f(x) \in \mathbb{F}_2[x]$ is irreducible, and thus $f(x)$ divides $F(x)$. Now, since the inversion $f^*(x) = x^n f(1/x)$ also belong to $\mathrm{PGL}_2(\mathbb{F}_q)$, it follows that $x^n f(1/x) = f(x)$ and so $f^*(x) = f(x)$ divides $F^*(x) = x^{2^j} + x^{2^j-1} + 1$. Thus $f(x)$ divides $F(x) + x(F^*(x) + F(x)) = x^2 + x + 1$. Since $\deg f(x) \geq 2$, the only possibility is $f(x) = x^2 + x + 1$. It can be easily verified that $f(x) = x^2 + x + 1 \in \mathbb{F}_2[x]$ is irreducible and $A \circ f = f$ for all $A \in \mathrm{PGL}_2(\mathbb{F}_2)$.

4. $p-$ subgroups of $\mathbf{GL_2(\mathbb{F}_q)}$

Let $p = \mathrm{char}\mathbb{F}_q$. If $S \subset \mathbb{F}_q$ is an $\mathbb{F}_p-$vector space of dimension r, then the set of translations $\{x + s; s \in S\}$ corresponds to the $p-$group H_S of the matrices $\begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}$ where s runs through S, $|H_S| = |S| = p^r$. This correspondence is an one-to-one correspondence between the subgroups of $H_{\mathbb{F}_q}$ and $\mathbb{F}_p-$ vector spaces of $\mathbb{F}_q$.

For a $p-$group $H \subset \mathrm{GL}_2(\mathbb{F}_q)$, $I_H(n)$ denotes the set of all monic irreducible polynomials in $\mathbb{F}_q$ of degree n such that $A \circ f = f$ for all $A \in H$. In the previous correspondence, the sets $I_{H_S}(n)$ and $C_S(n)$ (defined in Section 2) are the same.

Combining the observations above with Theorem 2 we are able to characterize the numbers $|I_H(n)|$:

Theorem 6. *Let $H \subset \mathrm{GL}_2(\mathbb{F}_q)$ be any group of order p^r.*

a) If $r > 1$ or n is not divisible by p, then $|I_H(n)| = 0$.

b) If $r = 1$ and $n = pm$, then

$$|I_H(n)| = \frac{p-1}{pm} \sum_{\substack{d|m \\ \gcd(d,p)=1}} q^{m/d}\mu(d).$$

Proof. Notice that $|\mathrm{GL}_2(\mathbb{F}_q)| = q(q-1)^2(q+1)$ and the sets of all translations in $\mathbb{F}_q$ corresponds to a Sylow $p-$ subgroup G of $\mathrm{GL}_2(\mathbb{F}_q)$. Let $H \subset \mathrm{GL}_2(\mathbb{F}_q)$ be any group of order p^r. We know that H is contained in some Sylow $p-$ *subgroup* K of $\mathrm{GL}_2(\mathbb{F}_q)$. Since all Sylow $p-$subgroups are conjugate, there is some $A \in \mathrm{GL}_2(\mathbb{F}_q)$ such that $A^{-1}KA = G$ and then $A^{-1}HA$ is a subgroup of G and has order p^r. Consider now the following map:

$$\begin{array}{rccc} \tau_{H,A}: & I_H(n) & \to & I_{A^{-1}HA}(n) \\ & f(x) & \mapsto & k_{A,f,n}(A^{-1} \circ f(x)), \end{array}$$

where $k_{A,f,n}$ is the only element in $\mathbb{F}_q$ such that $k_{A,f,n}(A^{-1} \circ f(x))$ is monic. A direct calculation shows that $\tau_{H,A}$ is well defined. Now, suppose that $\tau_{H,A}(f) = \tau_{H,A}(g)$ for some $f, g \in I_H(n)$, i.e., $k_{A,f,n}(A^{-1} \circ f(x)) = k_{A,g,n}(A^{-1} \circ g(x))$. Applying A we get $k_{A,f,n} f = k_{A,g,n} g$. Since f and g are monic irreducible we conclude that $f = g$ and then $\tau_{H,A}$ is one to one. In a similar way we can define a map from $I_{A^{-1}HA}(n)$ into $I_H(n)$. Thus $|I_{A^{-1}HA}(n)| = |I_H(n)|$.

The advantage is that $A^{-1}HA$ is a $p-$subgroup of G, i.e., $I_{A^{-1}HA}(n)$ is equal to $C_S(n)$ for some $\mathbb{F}_p-$vector space S of dimension r. Now, the result follows from Theorem 2. □

Acknowledgments

I would like to thank Daniel Panario for some helpful suggestions in the conclusion of this work and John MacQuarrie for the English corrections.

References

[1] S. Agou, *Irréductibilité des polynómes* $f(x^{p^{2r}} - ax^{p^r} - bx)$ *sur un corps fini* $\mathbb{F}_{p^s}$, Journal of Number Theory 10 (1979), pp 20.

[2] S. D. Cohen, *On irreducible polynomials of certain types in finite fields*, Mathematical Proceedings of the Cambridge Philosophical Society 66 (1969), pp 335-344.

[3] T. Garefalakis, *On the action of* $\mathrm{GL}_2(\mathbb{F}_q)$ *on irreducible polynomials over* $\mathbb{F}_q$, Journal of Pure and Applied Algebra. 215 (2011) pp 1835-1843.

[4] R. Lidl, H. Niederreiter, *Introduction to finite fields and their applications*. Cambridge University Press New York, NY, USA 1986.

[5] F. Ruskey, C. R. Miers, and J. Sawada, *The number of irreducible polynomials and Lyndon words with given trace*, SIAM J. Discrete Math. 14 (2001) pp 240-245.

[6] H. Stichtenoth, A. Topuzoglu, *Factorization of a class of polynomials over finite fields*, Finite Fields Appl. 18 (2012) pp 108-122.